

Pathwise uniqueness for a class of SPDEs driven by cylindrical α -stable processes ^{*}

Xiaobin Sun[†] Longjie Xie[‡] Yingchao Xie[§]

School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou 221116, China.

Abstract

We show the pathwise uniqueness for stochastic partial differential equation driven by a cylindrical α -stable process with Hölder continuous drift, thus obtaining an infinite dimensional generalization of the result of Priola [Osaka J. Math., 2012] in the case $H = \mathbb{R}^d$. The proof is based on an infinite dimensional Kolmogorov equation with non-local operator.

1 Introduction

In this paper, we consider the following stochastic partial differential equation (SPDE) in Hilbert space:

$$\begin{cases} dX_t = AX_t dt + B(X_t)dt + dZ_t, \\ X_0 = x \in H. \end{cases} \quad (1.1)$$

The objects are: a separable Hilbert space H with the inner product $\langle \cdot, \cdot \rangle$ and norm $|\cdot|$, a self-adjoint operator $A : \mathcal{D}(A) \subset H \rightarrow H$ which is the infinitesimal generator of a linear strongly continuous semigroup $(e^{tA})_{t \geq 0}$. The process $Z = (Z_t)_{t \geq 0}$ is a cylindrical α -stable process with $\alpha \in (1, 2)$ defined on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ equipped with the filtration $\{\mathcal{F}_t, t \geq 0\}$. We shall only assume $B : H \rightarrow H$ is Hölder continuous with certain power. Our aim is to prove the pathwise uniqueness for SPDE (1.1), which is an infinite dimensional generalization of the result by Priola [15] in the case $H = \mathbb{R}^d$.

Currently, there is an increasing interests in understanding the regularization effects of noise to the deterministic equations. We refer to [9] for a review on this direction. When $H = \mathbb{R}^d$ and $A = 0$, SPDE (1.1) is just the stochastic differential equation (SDE):

$$dX_t = b(X_t)dt + dZ_t, \quad X_0 = x \in \mathbb{R}^d. \quad (1.2)$$

In the case that Z_t is a Brownian motion, Veretennikov [23] first proved that SDE (1.2) has a unique global strong solution $X_t(x)$ if b is bounded and measurable. Later, Krylov and

^{*}Supported by the NNSF of China (No.: 11601196, 11271169) and the Priority Academic Program Development of Jiangsu Higher Education Institutions.

Keywords and phrases: Pathwise uniqueness; Stochastic partial differential equation; α -stable process

[†]E-mail: xbsun@jsnu.edu.cn

[‡]E-mail: xlj.98@whu.edu.cn

[§]E-mail: ycxie@jsnu.edu.cn

Röckner [11] shows that SDE (1.2) has a unique strong solution when $b \in L^p(\mathbb{R}^d)$ with $p > d$. The case that Z_t is a pure jump symmetric α -stable process has more difficulties. When $\alpha \geq 1$ and

$$b \in C_b^\beta(\mathbb{R}^d) \quad \text{with} \quad \beta > 1 - \frac{\alpha}{2},$$

it was proved by Priola [15] that there exists a unique strong solution $X_t(x)$ to SDE (1.2) for each $x \in \mathbb{R}^d$. Recently, Zhang [27] obtained the pathwise uniqueness to SDE (1.2) when $\alpha > 1$, b is bounded and belongs to certain fractional Sobolev space. See also [7, 8, 13, 16, 17, 28] and references therein for related results concerning SDEs with irregular coefficients.

For the infinite dimensional case, when Z_t is a cylindrical Wiener noise and B is Hölder continuous, the authors in [1] showed that there exists a unique strong solution to SPDE (1.1) for every $x \in H$. The extension of Veretennikov's result to infinite dimensional with B bounded was done in [2]. However, the uniqueness holds only for almost all starting point $x \in H$. See also [3, 4, 24, 25] and references therein.

Usually, SPDEs with jumps have more wide range of applications, we refer to the recent monograph [14]. When the drift B in (1.1) is Lipschitz continuous, the existence and uniqueness of solution can be easily obtained by a fixed point argument in [21]. Later on, more concreted SPDEs driven by cylindrical α -stable processes have been studied, see [5, 6, 19, 20, 21, 22, 26].

As far as we know, there is still no work on the pathwise uniqueness for SPDEs driven by pure jump Lévy process with irregular coefficient. The main difficult is that, from the analytic point of view, the generator of pure jump Lévy process is a non-local operator; and from the probability point of view, processes with jumps are more complicated than the continuous diffusion processes. We shall study the pathwise uniqueness of SPDEs (1.1) with Hölder continuous drift by solving the infinite dimensional Kolmogorov equation with non-local operator.

The paper proceeds as follows: In section 2, we state the main result. In Section 3, we study the regularity of the Ornstein-Uhlenbeck semigroup and solve the corresponding Kolmogorov equation. Finally, the proof of main result is given in Section 4.

Throughout our paper, we use the following convention: C with or without subscripts will denote a positive constant, whose value may change in different places, and whose dependence on parameters can be traced from calculations.

2 Preliminaries and main result

Given $\beta \in (0, 1]$, we denote by $C_b^\beta(H, H)$ the usual Hölder space of functions $G(x) : H \rightarrow H$ with norm

$$\|G\|_\beta := \sup_{x \in H} |G(x)| + \sup_{x \neq y \in H} \frac{|G(x) - G(y)|}{|x - y|^\beta},$$

and let $\|G\|_0 = \sup_{x \in H} |G(x)|$. Similar, for given $\beta \in (1, 2]$, the space $C_b^\beta(H, H)$ denotes functions satisfying

$$\|G\|_\beta := \|G\|_1 + \sup_{x \neq y \in H} \frac{\|DG(x) - DG(y)\|}{|x - y|^{\beta-1}} < \infty,$$

where $\|\cdot\|$ is the operator norm.

The cylindrical α -stable process Z is denoted by

$$Z_t = \sum_{n \geq 1} \beta_n Z_t^n e_n, \quad t \geq 0,$$

where $\{\beta_n\}_{n \geq 1}$ is a given sequence of positive numbers, $\{e_n\}_{n \geq 1}$ is a complete orthonormal basis of H , and $\{Z_t^n\}_{n \geq 1}$ are independent one dimensional rotationally symmetric α -stable process, i.e., the Lévy measure of Z_t^n is given by

$$\nu(dz) = \frac{c_\alpha}{|z|^{1+\alpha}} dz,$$

where c_α is a positive constant. By Lévy-Itô's decomposition, one has

$$Z_t^n = \int_{|x| \leq 1} x \tilde{N}^{(n)}(t, dx) + \int_{|x| > 1} x N^{(n)}(t, dx),$$

where $N^{(n)}(t, \Gamma)$ is the Poisson random measure, i.e.,

$$N^{(n)}(t, \Gamma) = \sum_{s \leq t} I_\Gamma(Z_s^n - Z_{s-}^n), \quad \forall t > 0, \Gamma \in \mathcal{B}(\mathbb{R} \setminus \{0\})$$

and $\tilde{N}^{(n)}(t, \Gamma)$ is compensated Poisson measure, i.e.,

$$\tilde{N}^{(n)}(t, \Gamma) = N^{(n)}(t, \Gamma) - t\nu(\Gamma).$$

Consider equation (1.1), we make the following assumptions:

- (i) $\{e_n\}_{n \geq 1} \subset \mathcal{D}(A)$, $Ae_n = -\gamma_n e_n$ with $\gamma_n > 0$ and $\gamma_n \uparrow \infty$.
- (ii) $\sum_{n \geq 1} \beta_n^\alpha < \infty$.
- (iii) $\sum_{n \geq 1} \frac{1}{\gamma_n} < \infty$.
- (iv) There exists a $\gamma \in (1, \alpha]$ such that for any $r < \gamma$ and $\lambda > 0$, we have

$$\int_0^\infty e^{-\lambda t} \Lambda_t^r dt < \infty, \tag{2.1}$$

where

$$\Lambda_t := \sup_{n \geq 1} \frac{e^{-\gamma_n t} \gamma_n^{1/\alpha}}{\beta_n} < \infty. \tag{2.2}$$

The main result of our paper is stated as follows:

Theorem 2.1 *Assume that (i)-(iv) hold and $B \in C_b^\beta(H, H)$ for some $\beta \in (1 + \alpha/2 - \gamma, 1)$. Then, SPDE (1.1) has a unique strong solution for each $x \in H$.*

Remark 2.2 The condition (ii) is the sufficient and necessary condition for Z_t is a Lévy process in H (see [12]). (2.1) and (2.2) in condition (iv) are used to study the smoothing property of the Ornstein-Uhlenbeck semigroup and regularity of the solution of the corresponding Kolmogorov equation in section 3.

We give some examples to illustrate our result.

Example 2.3 In the case that $H = \mathbb{R}^d$, we can take γ in (2.1) equals to α , and this means that we need $B \in C_b^\beta(\mathbb{R}^d)$ for $\beta \in (1 - \alpha/2, 1)$. Thus, we go back to [15].

Example 2.4 We set $H = L^2(D)$, where $D = [0, \pi]$, and denote by ∂D the boundary of D . Considering the stochastic Reaction-Diffusion Equation on D .

$$\begin{cases} dX(t, \xi) = \Delta_\xi^p X(t, \xi)dt + b(X(t, \xi))dt + dZ_t, \\ X(t, \xi) = 0, \quad t \geq 0, \quad \xi \in \partial D \\ X(0, \xi) = x(\xi), \quad \xi \in D, x \in H, \end{cases} \quad (2.3)$$

where Z is a cylindrical α -stable process with $\alpha \in (1, 2)$, Δ_ξ^p is pseudodifferential operator with $p \geq 1$ and $\frac{1}{2p} + 1 < \alpha$, $b : \mathbb{R} \rightarrow \mathbb{R}$ is a bounded and Hölder continuous with index β , where β will be determined later. Put

$$Ax = \Delta_\xi^p x, \quad x \in \mathcal{D}(A) = H^{2p}(D) \cap H_0^1(D),$$

where $H^{2p}(D)$ is the usual Sobolev space, and

$$B(x) = b(x(\cdot)), \quad x \in H.$$

Operator A possesses a complete orthonormal system of eigenfunctions namely

$$e_n(\xi) = (\sqrt{2/\pi}) \sin(n\xi), \quad \xi \in [0, \pi],$$

where $n \in \mathbb{N}$. The corresponding eigenvalues are $-\gamma_n$, where $\gamma_n = n^{2p}$.

Moreover, notice that $\frac{1}{2p} + 1 < \alpha$, if choosing $\beta_n = C\gamma_n^{-r}$ with $r \in \left(\frac{1}{2p\alpha}, \frac{\alpha-1}{\alpha}\right)$, then it is easy to verify conditions (i)-(iii) hold. Also by Remark 3.3 below, $\Lambda_t = \sup_{n \geq 1} \frac{e^{-\gamma_n t} \gamma_n^{1/\alpha}}{\beta_n} \leq \frac{C}{t^{r+\frac{1}{\alpha}}}$, and taking $\gamma = \frac{\alpha}{\alpha r + 1} \in (1, \alpha)$

$$\int_0^\infty e^{-\lambda t} \Lambda_t^q dt < \infty, \quad \forall q < \gamma,$$

which verifies condition (iv). Consequently, taking Hölder index $\beta \in \left(1 - \frac{\alpha(1-\alpha r)}{2(\alpha r + 1)}, 1\right)$, then (2.3) has a unique strong solution by Theorem 2.1.

For instance, if $p = 1$, i.e., Δ_ξ^p is the Laplace operator, then for any $\alpha \in (\frac{3}{2}, 2)$, taking

$$\beta_n = C\gamma_n^{-r} \quad \text{with} \quad r \in \left(\frac{1}{2\alpha}, \frac{\alpha-1}{\alpha}\right)$$

and

$$\gamma = \frac{\alpha}{\alpha r + 1} \in (1, \alpha), \quad \beta \in \left(1 - \frac{\alpha(1-\alpha r)}{2(\alpha r + 1)}, 1\right).$$

3 Ornstein-Uhlenbeck semigroup and corresponding Kolmogorov equation

3.1 H -valued Ornstein-Uhlenbeck semigroup

We first consider the following linear equation:

$$dY_t = AY_t dt + dZ_t, \quad Y_0 = x \in H. \quad (3.1)$$

Throughout this subsection, we assume that (i), (2.2) hold and

(ii)' There exists a positive constant C_α such that

$$\sum_{n=1}^{\infty} \frac{\beta_n^\alpha}{\gamma_n} \leq C_\alpha < \infty.$$

Notice that condition (ii)' is weaker than (ii). Under the assumptions (i) and (ii)', it is well-known that equation (3.1) has a unique mild solution for any initial value $x \in H$, which is given by

$$Y_t^x = e^{tA}x + Z_A(t),$$

where $Z_A(t) = \int_0^t e^{(t-s)A} dZ_s$. The solution Y_t is called the Ornstein-Uhlenbeck process and has received a lot of attentions. Let $R_t : B_b(H) \rightarrow B_b(H)$ be the corresponding semigroup defined by

$$R_t f(x) = \mathbb{E}[f(Y_t^x)] = \int_H f(y) \mu_t^x(dy), \quad x \in H, \quad f \in B_b(H), \quad t \geq 0,$$

where μ_t^x is the law of Y_t^x and $B_b(H)$ consists of all bounded functions $f : H \rightarrow \mathbb{R}$. This semigroup has been also studied under the name of generalized Mehler semigroup. It was shown in [21] that μ_t^x can be seen as Borel product measures in $\mathbb{R}^\infty$, i.e.,

$$\mu_t^x = \prod_{k \geq 1} \mu_t^{x_k},$$

where $x_k = \langle x, e_k \rangle$, and $\mu_t^{x_k}$ is a probability measure on $\mathbb{R}$ with density function

$$\frac{1}{c_k(t)} p_\alpha \left(\frac{z_k - e^{-\gamma_k t} x_k}{c_k(t)} \right), \quad \text{and} \quad c_k(t) := \beta_k \left(\frac{1 - e^{-\alpha \gamma_k t}}{\alpha \gamma_k} \right)^{1/\alpha}$$

here, p_α is the density of random variable Z_1^n .

The next result shows that R_t has a smoothing effect and the estimates of first and second derivative are given, which is an important step to prove our main result. Below, we also use $\langle \cdot, \cdot \rangle$ as the action of two elements without confusion.

Theorem 3.1 *For every $x, h, g \in H$ with $|h| \leq 1$, $|g| \leq 1$, set $h_k = \langle h, e_k \rangle$, $g_k = \langle g, e_k \rangle$. Then, for any $t > 0$ and $f \in B_b(H)$, we have $R_t f \in C_b^2(H)$ with*

(i) First order derivative:

$$\langle DR_t f(x), h \rangle = - \int_H f(z) \left(\sum_{k=1}^{\infty} \frac{p'_\alpha\left(\frac{z_k - e^{-\gamma_k t} x_k}{c_k(t)}\right)}{p_\alpha\left(\frac{z_k - e^{-\gamma_k t} x_k}{c_k(t)}\right)} \frac{e^{-\gamma_k t} h_k}{c_k(t)} \right) \mu_t^0(dz) \quad (3.2)$$

where μ_t^0 is the law of $Y_t^0 = Z_A(t)$, and

$$\sup_{x \in H} |\langle DR_t f(x), h \rangle| \leq c_\alpha \Lambda_t \|f\|_0, \quad \text{where} \quad c_\alpha = \int_{\mathbb{R}} \frac{p'_\alpha(z)^2}{p_\alpha(z)} dz, \quad (3.3)$$

and Λ_t is given by (2.2).

(ii) Second order derivative:

$$\begin{aligned} \langle D^2 R_t f(x) h, g \rangle &= - \int_H f(z + e^{tA} x) \sum_{l=1}^{\infty} \left(\sum_{k \neq l} \frac{p'_\alpha\left(\frac{z_k}{c_k(t)}\right) p'_\alpha\left(\frac{z_l}{c_l(t)}\right)}{p_\alpha\left(\frac{z_k}{c_k(t)}\right) p_\alpha\left(\frac{z_l}{c_l(t)}\right)} \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} \right. \\ &\quad \left. + \frac{p''_\alpha\left(\frac{z_l}{c_l(t)}\right)}{p_\alpha\left(\frac{z_l}{c_l(t)}\right)} \frac{e^{-2\gamma_l t} h_l g_l}{c_l^2(t)} \right) \mu_t^0(dz), \end{aligned} \quad (3.4)$$

and

$$\sup_{x \in H} |\langle D^2 R_t f(x) h, g \rangle| \leq \tilde{c}_\alpha \Lambda_t^2 \|f\|_0, \quad (3.5)$$

$$\text{where } \tilde{c}_\alpha = \sqrt{2} \max \left\{ \int_{\mathbb{R}} \frac{p'_\alpha(z)^2}{p_\alpha(z)} dz, \left[\int_{\mathbb{R}} \frac{p''_\alpha(z)^2}{p_\alpha(z)} dz \right]^{1/2} \right\}.$$

(iii) Hölder continuity: for any $\beta > 0$ and $r \in (0, 1)$,

$$|\langle DR_t f(x) - DR_t f(y), h \rangle| \leq C_\alpha \Lambda_t^{1+r-\beta} \|f\|_\beta |x - y|^r, \quad \forall x, y \in H, \quad (3.6)$$

where $C_\alpha > 0$ is a constant.

Proof The results that $R_t f \in C_b^1(H)$, (3.2), (3.3) have been proved in [21, Theorem 4.14], it suffices to prove (3.4)-(3.6). In order to show (3.4) and (3.5), we mainly follow the steps in [21, Theorem 4.14]. So, we only consider the case that $f \in C_b(H)$ is cylindrical, i.e.,

$$f(x) = \tilde{f}(x_1, \dots, x_j), \quad x \in H$$

for some $j \geq 1$ and $\tilde{f} : \mathbb{R}^j \rightarrow \mathbb{R}$. We also assume that $\tilde{f}$ has bounded support in $\mathbb{R}^j$. Then the general case of f can be proved by an argument of approximation (see Step II-Step V in [21, Theorem 4.14]).

Fix arbitrary $x, h, g \in H$ with $|h| \leq 1, |g| \leq 1$. we will show that there exists $D_{h \otimes g}^2 R_t f(x)$, the directional derivative of $DR_t f(x)h$, along the direction g at x . To shorten the notation, we write

$$\xi_{k,l}(z) := \frac{p'_\alpha\left(\frac{z_k - e^{-\gamma_k t} x_k}{c_k(t)}\right) p'_\alpha\left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)}\right)}{p_\alpha\left(\frac{z_k - e^{-\gamma_k t} x_k}{c_k(t)}\right) p_\alpha\left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)}\right)}.$$

Let $g^N = \sum_{k=1}^N g_k e_k$, for any $m \geq \max\{j, N\}$. Then by (3.2), we get

$$\begin{aligned} \langle D^2 R_t f(x) h, g^N \rangle &= - \int_{\mathbb{R}^m} \tilde{f}(z) \sum_{l=1}^N \left(\sum_{k \neq l} \xi_{k,l}(z) \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} \right. \\ &\quad \left. + \frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) (e^{-\gamma_l t})^2 h_l g_l}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) c_l^2(t)} \right) \prod_{l=1}^m \mu_t^{x_l}(dz_l) \\ &= - \int_H f(z) \sum_{l=1}^N \left(\sum_{k \neq l} \xi_{k,l}(z) \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} \right. \\ &\quad \left. + \frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) e^{-2\gamma_l t} h_l g_l}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) c_l^2(t)} \right) \mu_t^x(dz). \end{aligned}$$

In order to pass to the limit, as $N \rightarrow \infty$, we show that

$$\begin{aligned} \phi_N(t, x) &:= \sum_{l=1}^N \left(\sum_{k \neq l} \xi_{k,l}(z) \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} \right. \\ &\quad \left. + \frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) e^{-2\gamma_l t} h_l g_l}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) c_l^2(t)} \right) \text{ converges in } L^2(\mu_t^x). \end{aligned} \quad (3.7)$$

In fact, notice that for any $k \neq l$,

$$\begin{aligned} &\int_H \xi_{k,l}(z) \cdot \frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)} \mu_t^x(dz) \\ &= \int_{\mathbb{R}} \frac{p'_{\alpha} \left(\frac{z_k - e^{-\gamma_k t} x_k}{c_k(t)} \right)}{p_{\alpha} \left(\frac{z_k - e^{-\gamma_k t} x_k}{c_k(t)} \right)} \mu_t^{x_k}(dz_k) \int_{\mathbb{R}} \frac{p'_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)} \cdot \frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)} \mu_t^{x_l}(dz_l) \\ &= \int_{\mathbb{R}} p'_{\alpha}(y) dy \cdot \int_{\mathbb{R}} \frac{p'_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)} \cdot \frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right)} \mu_t^{x_l}(dz_l) \\ &= 0, \end{aligned}$$

where the last inequality by the fact that p'_{α} is odd. Then, for any $N, p \in \mathbb{N}$,

$$\begin{aligned} &\int_H \left| \sum_{l=N}^{N+p} \left(\sum_{k \neq l} \xi_{k,l}(z) \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} + \frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) (e^{-\gamma_l t})^2 h_l g_l}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) c_l^2(t)} \right) \right|^2 \mu_t^x(dz) \\ &= \int_H \sum_{l=N}^{N+p} \sum_{k \neq l} \left(\xi_{k,l}(z) \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} \right)^2 + \sum_{l=N}^{N+p} \left(\frac{p''_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) (e^{-\gamma_l t})^2 h_l g_l}{p_{\alpha} \left(\frac{z_l - e^{-\gamma_l t} x_l}{c_l(t)} \right) c_l^2(t)} \right)^2 \mu_t^x(dz) \\ &= \sum_{l=N}^{N+p} \sum_{k \neq l} \left(\frac{e^{-\gamma_k t} e^{-\gamma_l t}}{c_k(t) c_l(t)} \right)^2 h_k^2 g_l^2 \int_{\mathbb{R}} \frac{p'_{\alpha}(y_l)^2}{p_{\alpha}(y_l)} dy_l \int_{\mathbb{R}} \frac{p'_{\alpha}(y_k)^2}{p_{\alpha}(y_k)} dy_k + \sum_{l=N}^{N+p} \left(\frac{e^{-\gamma_l t}}{c_l^2(t)} \right)^4 h_l^2 g_l^2 \int_{\mathbb{R}} \frac{p''_{\alpha}(y_l)}{p_{\alpha}(y_l)} dy_l \\ &\leq \tilde{c}_{\alpha}^2 \Lambda_t^4 |h|^2 \sum_{l=N}^{N+p} g_l^2, \end{aligned}$$

where $\tilde{c}_\alpha = \sqrt{2} \max \left\{ \int_{\mathbb{R}} \frac{p'_\alpha(z)^2}{p_\alpha(z)} dz, \left[\int_{\mathbb{R}} \frac{p''_\alpha(z)^2}{p_\alpha(z)} dz \right]^{1/2} \right\}$.

Note that, for any $N \in \mathbb{N}$,

$$\begin{aligned} \langle D^2 R_t f(x) h, g^N \rangle &= - \int_H f(z + e^{tA} x) \sum_{l=1}^N \left(\sum_{k \neq l} \frac{p'_\alpha(\frac{z_k}{c_k(t)}) p'_\alpha(\frac{z_l}{c_l(t)})}{p_\alpha(\frac{z_k}{c_k(t)}) p_\alpha(\frac{z_l}{c_l(t)})} \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} \right. \\ &\quad \left. + \frac{p''_\alpha(\frac{z_l}{c_l(t)})}{p_\alpha(\frac{z_l}{c_l(t)})} \frac{(e^{-\gamma_l t})^2 h_l g_l}{c_l^2(t)} \right) \mu_t^0(dz). \end{aligned}$$

Up to now we can showed that

$$\frac{DR_t f(x + \epsilon g^N) h - DR_t f(x) h}{\epsilon} = \frac{1}{\epsilon} \int_0^\epsilon \langle D^2 R_t f(x + r g^N) h, g^N \rangle dr \quad (3.8)$$

Using (3.7), it is easy to see that, for any $r \in (0, 1)$, $N \in \mathbb{N}$,

$$\begin{aligned} &\lim_{N \rightarrow \infty} \langle D^2 R_t f(x + r g^N) h, g^N \rangle \\ &= - \int_H f(z + e^{tA}(x + r g)) \sum_{l=1}^\infty \left(\sum_{k \neq l} \frac{p'_\alpha(\frac{z_k}{c_k(t)}) p'_\alpha(\frac{z_l}{c_l(t)})}{p_\alpha(\frac{z_k}{c_k(t)}) p_\alpha(\frac{z_l}{c_l(t)})} \frac{e^{-\gamma_k t} e^{-\gamma_l t} h_k g_l}{c_k(t) c_l(t)} \right. \\ &\quad \left. + \frac{p''_\alpha(\frac{z_l}{c_l(t)})}{p_\alpha(\frac{z_l}{c_l(t)})} \frac{(e^{-\gamma_l t})^2 h_l g_l}{c_l^2(t)} \right) \mu_t^0(dz). \end{aligned} \quad (3.9)$$

Moreover, for any $r \in (0, 1)$, $|\langle D^2 R_t f(x + r g^N) h, g^N \rangle| \leq \tilde{c}_\alpha \Lambda_t^2 \|f\|_0$. Then by dominated convergence theorem in (3.8), we obtain

$$\frac{DR_t f(x + \epsilon g) h - DR_t f(x) h}{\epsilon} = \frac{1}{\epsilon} \int_0^\epsilon u(t, h, x + r g) dr, \quad \epsilon \in (0, 1),$$

where $u(t, x + r g)$ is the right-hand side of (3.9). This shows that $DR_t f(x) h$ is Gâteaux differentiable at $x \in H$ along the direction h and (3.4), (3.5) hold.

We proceed to show (3.6). It is easy to see that

$$\sup_{x \in H} |\langle DR_t f(x), h \rangle| \leq \|f\|_1. \quad (3.10)$$

By taking ∇f into (3.3), we can also obtain

$$\sup_{x \in H} |\langle D^2 R_t f(x) h, g \rangle| \leq c_\alpha \Lambda_t \|f\|_1. \quad (3.11)$$

Thus, we have by (3.3) and (3.10) that

$$\sup_{x \in H} |\langle DR_t f(x), h \rangle| \leq \hat{c}_\alpha \Lambda_t^{1-\beta} \|f\|_\beta,$$

and similarly, it holds by (3.5) and (3.11) that

$$\sup_{x \in H} |\langle D^2 R_t f(x) h, g \rangle| \leq \hat{c}_\alpha \Lambda_t^{2-\beta} \|f\|_\beta.$$

Combing the above two inequalities with the interpolation theory, we can get the desired result. ■

Remark 3.2 By the similar argument above, we could obtain $R_t f \in C_b^\infty(H)$, for any $t > 0$, $f \in B_b(H)$. Moreover, there exists a positive constant C_n such that

$$\sup_{x \in H} \|D^n R_t f(x)\| \leq C_n \Lambda_t^n \|f\|_0.$$

Remark 3.3 If there exists $r \geq -1/\alpha$ such that

$$\beta_n \geq C \gamma_n^{-r}, \quad \forall n \geq 1,$$

then we can give a upper bound of Λ_t in Theorem 3.1, i.e.,

$$\Lambda_t = \sup_{n \geq 1} \frac{e^{-\gamma_n t} \gamma_n^{1/\alpha}}{\beta_n} \leq \frac{C}{t^{r+\frac{1}{\alpha}}}, \quad t > 0.$$

3.2 Elliptic Kolmogorov equation

In this subsection, we always assume that (i), (ii)' and (iv) hold. Now, we study the following Kolmogorov equation in H

$$\lambda U - \langle B, DU \rangle - \mathcal{L}U = F, \quad (3.12)$$

where $\lambda > 0$, $F \in B_b(H, H)$ and the operator $\mathcal{L}$ is the infinitesimal generator of OU-semigroup R_t , i.e.,

$$\mathcal{L}U(x) = \langle Ax, DU(x) \rangle + \sum_{k=1}^{\infty} \beta_k^\alpha \int_{\mathbb{R}} [U(x + e_k z) - U(x) - \langle DU(x), e_k z \rangle 1_{\{|z| \leq 1\}}] \frac{c_\alpha}{|z|^{1+\alpha}} dz.$$

Theorem 3.4 Assume that $B \in C_b^\beta(H, H)$ for some $\beta > 0$. Then, for λ big enough, every $F \in C_b^\beta(H, H)$ and any $0 < \theta < \beta$, there exists a function $U \in C_b^{\gamma+\theta}(H, H)$ satisfying the following integral equation:

$$U(x) = \int_0^\infty e^{-\lambda t} R_t \left(\langle B, DU \rangle + F \right) (x) dt. \quad (3.13)$$

Moreover, U also solves equation (3.12) and we have

$$\|U\|_{\gamma+\theta} \leq C_\lambda \|F\|_\beta, \quad (3.14)$$

where C_λ is a positive constant satisfying $\lim_{\lambda \rightarrow +\infty} C_\lambda = 0$.

Proof Let us first show the estimate (3.14). In fact, let U satisfies (3.13), without loss of generality, we may assume that $1 + \beta < \gamma + \theta < 2$. Then, by (3.6) and condition (v) we have

$$\begin{aligned} \|U\|_{\gamma+\theta} &\leq \int_0^\infty e^{-\lambda t} \left\| R_t \left(\langle B, DU \rangle + F \right) \right\|_{\gamma+\theta} dt \\ &\leq \int_0^\infty e^{-\lambda t} \Lambda_t^{\gamma+\theta-\beta} dt \cdot \|\langle B, DU \rangle + F\|_\beta \\ &\leq C_\lambda \left(\|F\|_\beta + \|B\|_\beta \cdot \|U\|_{1+\beta} \right), \end{aligned}$$

where C_λ is given by

$$C_\lambda := \int_0^\infty e^{-\lambda t} \Lambda_t^{\gamma+\theta-\beta} dt,$$

and by dominate convergence theorem it holds that $\lim_{\lambda \rightarrow +\infty} C_\lambda = 0$. Take λ big enough such that

$$C_\lambda \|B\|_\beta < \frac{1}{2},$$

we get the desired result.

Now, we construct the solution of (3.13) via Picard's iteration argument. Set $U_0 \equiv 0$ and for $n \in \mathbb{N}$, define U_n recursively by

$$U_n(x) := \int_0^\infty e^{-\lambda t} R_t \left(\langle B, DU_{n-1} \rangle + F \right) (x) dt.$$

In view of (3.3), it is easy to check that $U_1 \in C_b^1(H, H)$, and U_2 is thus well defined, and so on. We show that $U_1 \in C_b^{\gamma+\theta}(H, H)$ with any $\theta \in (0, \beta)$. In fact, thanks to (v) and using (3.6) once again, we can deduce that

$$\begin{aligned} DU_1(x) - DU_1(y) &= \int_0^\infty e^{-\lambda t} \left[DR_t F(x) - DR_t F(y) \right] dt \\ &\leq c_\alpha |x - y|^{\gamma+\theta-1} \int_0^\infty e^{-\lambda t} \Lambda_t^{\gamma+\theta-\beta} dt \cdot \|F\|_\beta = c_\alpha C_\lambda |x - y|^{\gamma+\theta-1} \|F\|_\beta. \end{aligned}$$

Notice that $\gamma + \theta - 1 > \beta$. As a result,

$$\langle B, DU_1 \rangle \in C_b^\beta(H, H).$$

Repeating the above argument, we have for every $n \in \mathbb{N}$ and any $\theta \in (0, \beta)$,

$$U_n \in C_b^{\gamma+\theta}(H, H).$$

Moreover, for any $n > m$

$$U_n(x) - U_m(x) = \int_0^\infty e^{-\lambda t} R_t \left(\langle B, DU_{n-1} - DU_{m-1} \rangle \right) (x) dt,$$

we further have that

$$\begin{aligned} \|U_n - U_m\|_{\gamma+\theta} &\leq \int_0^\infty e^{-\lambda t} \left\| R_t \left(\langle B, DU_{n-1} - DU_{m-1} \rangle \right) \right\|_{\gamma+\theta} dt \\ &\leq c_\alpha \int_0^\infty e^{-\lambda t} \Lambda_t^{\gamma+\theta-\beta} dt \cdot \|\langle B, DU_{n-1} - DU_{m-1} \rangle\|_\beta \\ &\leq 2c_\alpha C_\lambda \|B\|_\beta \cdot \|DU_{n-1} - DU_{m-1}\|_\beta \\ &\leq 2c_\alpha C_\lambda \|B\|_\beta \cdot \|U_{n-1} - U_{m-1}\|_{\gamma+\theta}, \end{aligned}$$

This means that for λ big enough, U_n is Cauchy sequence in $C_b^{\gamma+\theta}(H, H)$. Thus, there exists a limit function $U \in C_b^{\gamma+\theta}(H, H)$ with $\theta \in (0, \beta)$ satisfying (3.13). The assertion that U solves (3.12) follows by integral by part. The whole proof is finished. ■

Remark 3.5 *It may be expected that the optimal regularity of U should be $C_b^{\alpha+\beta}(H, H)$. However, due to the uncertain of Λ_t , we can not obtain thus estimate. Nevertheless, this is enough for us to prove the pathwise uniqueness for SPDE (1.1).*

4 Strong uniqueness

We call a predictable H -valued stochastic process $\{X_t\}_{t \geq 0}$ depending on initial value $x \in H$, is a mild solution of equation (1.1) on the filtered probability space $(\Omega, \mathcal{F}_t, \mathbb{P})$, if $\{X_t\}_{t \geq 0}$ is an $\mathcal{F}_t$ -adapted and satisfies

$$X_t = e^{tA}x + \int_0^t e^{(t-s)A}B(X_s)ds + \int_0^t e^{(t-s)A}dZ_s, \quad (4.1)$$

where the deterministic integral in (4.1) is well defined by the assumption that B is bounded. The existence of a solution to equation (1.1) is known under our assumptions. Thus, by the classical Yamada-Watanabe principle [10], we only need to focus on the pathwise uniqueness.

Usually, the Itô's formula is performed for functions $f \in C_b^2(H, H)$. However, this is too strong for our latter use. Actually, Z_t is a α -stable process in our case, and we will show that Itô's formula holds for $f \in C_b^r(H, H)$ with any $r > \alpha$, which is stated as follows:

Lemma 4.1 *Let X_t satisfies equation (1.1) and $f \in C_b^r(H, H)$ with $r > \alpha$. Then, we have*

$$\begin{aligned} f(X_t) &= f(x) + \int_0^t \langle AX_s + B(X_s), Df(X_s) \rangle ds \\ &\quad + \sum_{k=1} \beta_k^\alpha \int_0^t \int_{\mathbb{R}} [f(X_s + e_k z) - f(X_s) - \langle Df(X_s), e_k z \rangle 1_{\{|z| \leq 1\}}] \frac{1}{|z|^{1+\alpha}} dz ds \\ &\quad + \sum_{k=1} \int_0^t \int_{\mathbb{R}} [f(X_{s-} + \beta_k e_k z) - f(X_{s-})] \tilde{N}^{(k)}(dz, ds). \end{aligned}$$

Proof Let $f_n \in C_b^2(H, H)$ with $\|f_n\|_r \leq \|f\|_r$ and $\|f_n - f\|_{r'} \rightarrow 0$ for every $r' < r$. Then we can use Itô's formula for $f_n(X_t)$, i.e.,

$$\begin{aligned} f_n(X_t) &= f_n(x) + \int_0^t \langle AX_s + B(X_s), Df_n(X_s) \rangle ds \\ &\quad + \sum_{k=1} \beta_k^\alpha \int_0^t \int_{\mathbb{R}} [f_n(X_s + e_k z) - f_n(X_s) - \langle Df_n(X_s), e_k z \rangle 1_{\{|z| \leq 1\}}] \frac{1}{|z|^{1+\alpha}} dz ds \\ &\quad + \sum_{k=1} \int_0^t \int_{\mathbb{R}} [f_n(X_{s-} + \beta_k e_k z) - f_n(X_{s-})] \tilde{N}^{(k)}(dz, ds) \\ &=: \mathbb{I}_1 + \mathbb{I}_2 + \mathbb{I}_3. \end{aligned}$$

Now we are going to pass the limits on the both sides of the above equality. For any $x \in H$, it is easy to see that, as $n \rightarrow \infty$,

$$\mathbb{I}_1 \rightarrow f(x) + \int_0^t \langle AX_s + B(X_s), Df(X_s) \rangle ds.$$

Thanks to the assumption that $\sum_{n \geq 1} \beta_n^\alpha < \infty$ and in view of the following estimates:

$$|f_n(x + e_k z) - f_n(x) - \langle Df_n(x), e_k z \rangle| \leq \|f_n\|_r |z|^r \leq \|f\|_r |z|^r, \quad |z| \leq 1$$

and

$$|f_n(x + e_k z) - f_n(x)| \leq 2\|f_n\|_0 \leq 2\|f\|_0, \quad |z| \geq 1,$$

we obtain by dominated convergence theorem that, as $n \rightarrow \infty$,

$$\mathbb{I}_2 \rightarrow \sum_{k=1} \beta_k^\alpha \int_0^t \int_{\mathbb{R}} [f(X_s + e_k z) - f(X_s) - \langle Df(X_s), e_k z \rangle 1_{\{|z| \leq 1\}}] \frac{1}{|z|^{1+\alpha}} dz ds.$$

Finally, by the isometry formula we have

$$\begin{aligned} & \mathbb{E} \left| \sum_{k=1} \int_0^t \int_{\mathbb{R}} [f_n(X_{s-} + \beta_k e_k z) - f_n(X_{s-}) - f(X_{s-} + \beta_k e_k z) + f(X_{s-})] \tilde{N}^{(k)}(dz, ds) \right|^2 \\ &= \sum_{k=1} \int_0^t \int_{\mathbb{R}} \mathbb{E} |f_n(X_s + \beta_k e_k z) - f_n(X_s) - f(X_{s-} + \beta_k e_k z) + f(X_{s-})|^2 \nu(dz) ds \\ &= \sum_{k=1} \beta_k^\alpha \int_0^t \int_{\mathbb{R}} \mathbb{E} |f_n(X_s + e_k z) - f_n(X_s) - f(X_s + e_k z) + f(X_s)|^2 \frac{1}{|z|^{1+\alpha}} dz ds \\ &\rightarrow 0, \quad n \rightarrow \infty. \end{aligned}$$

The proof is finished. ■

Now, assume $B \in C_b^\beta(H, H)$ and let U solves the following equation:

$$\lambda U - \langle B, DU \rangle - \mathcal{L}U = B.$$

According to Theorem 3.4, we have $U \in C_b^{\gamma+\theta}(H, H)$ with $\theta < \beta$. We prove the following Zvonkin's transformation.

Lemma 4.2 *Let X_t be a solution of equation 1.1, then we have*

$$\begin{aligned} X_t &= e^{tA}(x + U(x)) + \int_0^t e^{(t-s)A} \lambda U(X_s) ds - U(X_t) - \int_0^t A e^{(t-s)A} U(X_s) ds \\ &\quad + \sum_{k=1} \int_0^t \int_{\mathbb{R}} e^{(t-s)A} [U(X_{s-} + \beta_k e_k z) - U(X_{s-})] \tilde{N}^{(k)}(ds, dz) + \int_0^t e^{(t-s)A} dZ_s. \end{aligned} \quad (4.2)$$

Proof Thanks to Lemma 4.1, we can use the Itô's formula for $U(X_t)$ to get that

$$\begin{aligned} dU(X_t) &= \langle B(X_t), DU(X_t) \rangle dt + \mathcal{L}U(X_t) dt \\ &\quad + \sum_{k=1} \int_{\mathbb{R}} [U(X_{t-} + \beta_k e_k z) - U(X_{t-})] \tilde{N}^{(k)}(t, dz) \\ &= \lambda U(X_t) dt - B(X_t) dt + \sum_{k=1} \int_{\mathbb{R}} [U(X_{t-} + \beta_k e_k z) - U(X_{t-})] \tilde{N}^{(k)}(dt, dz), \end{aligned}$$

which give a formula for $B(X_t)dt$:

$$B(X_t)dt = \lambda U(X_t)dt - dU(X_t) + \sum_{k=1} \int_{\mathbb{R}} [U(X_{t-} + \beta_k e_k z) - U(X_{t-})] \tilde{N}^{(k)}(dt, dz).$$

We put this formula in equation (1.1) and get

$$\begin{aligned} dX_t &= AX_t dt + \lambda U(X_t) dt - dU(X_t) \\ &\quad + \sum_{k=1} \int_{\mathbb{R}} [U(X_{t-} + \beta_k e_k z) - U(X_{t-})] \tilde{N}^{(k)}(dt, dz) + dZ_t, \end{aligned}$$

then follow the usual variation of constant method and get

$$\begin{aligned} X_t &= e^{tA} x + \int_0^t e^{(t-s)A} \lambda U(X_s) ds - \int_0^t e^{(t-s)A} dU(X_s) \\ &\quad + \sum_{k=1} \int_0^t \int_{\mathbb{R}} e^{(t-s)A} [U(X_{s-} + \beta_k e_k z) - U(X_{s-})] \tilde{N}^{(k)}(ds, dz) + \int_0^t e^{(t-s)A} dZ_s. \end{aligned}$$

Finally, the integration by parts formula implies

$$\begin{aligned} X_t &= e^{tA}(x + U(x)) + \int_0^t e^{(t-s)A} \lambda U(X_s) ds - U(X_t) - \int_0^t A e^{(t-s)A} U(X_s) ds \\ &\quad + \sum_{k=1} \int_0^t \int_{\mathbb{R}} e^{(t-s)A} [U(X_{s-} + \beta_k e_k z) - U(X_{s-})] \tilde{N}^{(k)}(ds, dz) + \int_0^t e^{(t-s)A} dZ_s. \end{aligned}$$

The proof is complete. ■

The following result was proved in [15] in finite dimensional. For the sake of completeness, we provide a simple proof here.

Lemma 4.3 *For every $x, y \in H$ and any $\theta < \beta$, we have*

$$|U(x + z) - U(x) - U(y + z) + U(y)| \leq |x - y| \cdot |z|^{\gamma+\theta-1} \|U\|_{\gamma+\theta}. \quad (4.3)$$

Proof Set

$$J_z U(x) := U(x + z) - U(x).$$

It is obvious that for any $|z| \leq 1$,

$$\|DJ_z U\|_0 \leq |z|^{\gamma+\theta-1} \|DU\|_{\gamma+\theta-1}.$$

Thus, we can deduce that

$$|J_z U(x) - J_z U(y)| \leq |x - y| \cdot \|DJ_z U\|_0 \leq |x - y| \cdot |z|^{\gamma+\theta-1} \|U\|_{\gamma+\theta}.$$

The proof is complete. ■

We are now in the position to give the proof of our main result.

Proof of Theorem 2.1: Since uniqueness is a local property, we only need to consider on the interval $[0, T]$ with T small. Let X_t and Y_t be two solutions of equation (1.1) both starting from $x \in H$. Then, by (4.2) we have $V_t := X_t - Y_t$ satisfies the following equation:

$$V_t = \lambda \int_0^t e^{(t-s)A} (U(X_s) - U(Y_s)) ds - [U(X_t) - U(Y_t)]$$

$$- \int_0^t A e^{(t-s)A} \left(U(X_s) - U(Y_s) \right) ds + I_t(X) - I_t(Y),$$

where

$$I_t(X) = \sum_{k=1} \int_0^t \int_{\mathbb{R}} e^{(t-s)A} \left[U(X_{s-} + \beta_k e_k z) - U(X_{s-}) \right] \tilde{N}^{(k)}(ds, dz)$$

and

$$I_t(Y) = \sum_{k=1} \int_0^t \int_{\mathbb{R}} e^{(t-s)A} \left[U(Y_{s-} + \beta_k e_k z) - U(Y_{s-}) \right] \tilde{N}^{(k)}(ds, dz).$$

In view of (3.14), we have

$$\|U\|_{\gamma+\theta} \leq C_\lambda \|B\|_\beta.$$

Notice that by the maximal inequality,

$$\left\| \int_0^\cdot A e^{(\cdot-s)A} f_s ds \right\|_{L^2(0,T;H)}^2 \leq C_T \|f\|_{L^2(0,T;H)}^2,$$

where C_T is a constant independent of f . Then we have the following estimate:

$$\begin{aligned} \int_0^T |V_t|^2 dt &\leq 4 \int_0^T \left| \lambda \int_0^t e^{(t-s)A} \left(U(X_s) - U(Y_s) \right) ds \right|^2 dt \\ &\quad + 4 \int_0^T |U(X_t) - U(Y_t)|^2 dt + 4 \int_0^T \left| \int_0^t A e^{(t-s)A} \left(U(X_s) - U(Y_s) \right) ds \right|^2 dt \\ &\quad + 4 \int_0^T |I_t(X) - I_t(Y)|^2 dt \\ &\leq (4\lambda^2 T C_\lambda^2 \|B\|_\beta^2 + 4C_\lambda^2 \|B\|_\beta^2 + 4C_T C_\lambda^2 \|B\|_\beta^2) \int_0^T |V_t|^2 dt \\ &\quad + 4 \int_0^T |I_t(X) - I_t(Y)|^2 dt. \end{aligned}$$

Since $\lim_{\lambda \rightarrow \infty} C_\lambda = 0$, taking λ sufficient large such that $4C_\lambda^2 \|B\|_\beta^2 + 4C_T C_\lambda^2 \|B\|_\beta^2 \leq 1/3$ and T small enough such that $4\lambda^2 T C_\lambda^2 \|B\|_\beta^2 \leq 1/3$, then we have

$$\int_0^T |V_t|^2 dt \leq 12 \int_0^T |I_t(X) - I_t(Y)|^2 dt.$$

Meanwhile,

$$\begin{aligned} &\mathbb{E}|I_t(X) - I_t(Y)|^2 \\ &= \sum_{k=1} \int_0^t \int_{\mathbb{R}} \mathbb{E} \left| e^{(t-s)A} \left[U(X_{s-} + \beta_k e_k z) - U(X_s) - U(Y_s + \beta_k e_k z) + U(Y_{s-}) \right] \right|^2 \nu(dz) ds \\ &\leq \sum_{k=1} \beta_k^\alpha \int_0^t \sum_{m=1} e^{-2(t-s)\gamma_m} \int_{|z| \leq 1} \mathbb{E}|X_s - Y_s|^2 \cdot |z|^{2(\gamma+\theta-1)} \|U\|_{\gamma+\theta}^2 \nu(dz) ds \\ &\quad + \sum_{k=1} \beta_k^\alpha \int_0^t \sum_{m=1} e^{-2(t-s)\gamma_m} \int_{|z| > 1} 2 \|DU\|_0^2 \mathbb{E}|X_s - Y_s|^2 \nu(dz) ds \end{aligned}$$

Since we assumed that $\beta \in (1 + \alpha/2 - \gamma, 1)$, there always exists a $\theta < \beta$ such that

$$2(\gamma + \theta - 1) > \alpha.$$

As a result, we have by (4.3)

$$\mathbb{E}|I_t(X) - I_t(Y)|^2 \leq C_\lambda \|B\|_\beta^2 \sum_{k=1} \beta_k^\alpha \int_0^t \sum_{m=1} e^{-2(t-s)\gamma_m} \mathbb{E}|V_s|^2 ds.$$

Hence,

$$\begin{aligned} \mathbb{E} \int_0^T |I_t(X) - I_t(Y)|^2 dt &\leq C_\lambda \|B\|_\beta^2 \sum_{k=1} \beta_k^\alpha \int_0^T \int_0^t \sum_{m=1} e^{-2(t-s)\gamma_m} \mathbb{E}|V_s|^2 ds dt \\ &\leq C_\lambda \|B\|_\beta^2 \sum_{k=1} \beta_k^\alpha \int_0^T \left(\int_s^T \sum_{m=1} e^{-2(t-s)\gamma_m} dt \right) \mathbb{E}|V_s|^2 ds \\ &\leq C_\lambda \|B\|_\beta^2 \sum_{k=1} \beta_k^\alpha \left(\int_0^T \sum_{m=1} e^{-2t\gamma_m} dt \right) \int_0^T \mathbb{E}|V_s|^2 ds \\ &\leq C_\lambda \|B\|_\beta^2 \sum_{k=1} \beta_k^\alpha \sum_{m=1} \frac{1 - e^{-2T\gamma_m}}{2\gamma_m} \int_0^T \mathbb{E}|V_s|^2 ds \end{aligned}$$

By assumption, $\sum_{m=1} \frac{1 - e^{-2T\gamma_m}}{2\gamma_m}$ is finite and when $T \rightarrow 0$, it converges to zero. Therefore we can obtain $\mathbb{E} \int_0^T |I_t(X) - I_t(Y)|^2 dt = 0$ as T small enough. This implies $X = Y$. The proof is complete.

References

- [1] G. Da Prato and F. Flandoli. *Pathwise uniqueness for a class of SPDEs in Hilbert spaces and applications*. J. Funct. Anal., 259, 243–267, 2010.
- [2] G. Da Prato, F. Flandoli, E. Priola and M. Röckner. *Strong uniqueness for stochastic evolution equations in Hilbert spaces perturbed by a bounded measurable drift*. Ann. Probab., 41, 3306–3344, 2013.
- [3] G. Da Prato, F. Flandoli, E. Priola and M. Röckner. *Strong Uniqueness for Stochastic Evolution Equations with Unbounded Measurable Drift Term*. J. Theor. Prob., 28 (4), 1571–1600, 2015.
- [4] G. Da Prato, F. Flandoli, M. Röckner and A. Yu. Veretennikov. *Strong uniqueness for SDEs in Hilbert spaces with non-regular drift*. Ann. Probab. 44 (3), 1985–2023, 2016.
- [5] Z. Dong, L. Xu, and X. Zhang. *Invariance measures of stochastic 2D Navier-stokes equations driven by α -stable processes*. Electron. Commun. Probab., 16, 678–688, 2011.
- [6] Z. Dong, L. Xu, and X. Zhang. *Exponential ergodicity of stochastic Burgers equations driven by α -stable processes*. J. Stat. Phys., 4, 929–949, 2014.

- [7] E. Fedrizzi and F. Flandoli. *Hölder Flow and Differentiability for SDEs with Nonregular Drift*. Stoch. Anal. Appl., 31, 708–736, 2013.
- [8] E. Fedrizzi and F. Flandoli. *Noise prevents singularities in linear transport equations*. J. Funct. Anal., 264, 1329–1354, 2013.
- [9] F. Flandoli. *Random perturbation of PDEs and fluid dynamic models*. Lecture Notes in Mathematics. Springer, Heidelberg, 2011.
- [10] N. Ikeda and S. Watanabe. *Stochastic Differential Equations and Diffusion Processes, 2nd edition*. North-Holland, Kodansha, 1989.
- [11] N. V. Krylov and M. Röckner. *Strong solutions of stochastic equations with singular time dependent drift*. Probab. Theory Related Fields, 131 (2), 154–196, 2005.
- [12] Y. Liu and J. Zhai. *A note on time regularity of generalized Ornstein-Uhlenbeck processes with cylindrical stable noise*. C. R. Math. Acad. Sci. Paris, 350, 1–2, 2012.
- [13] O. Menoukeu-Pamen, T. Meyer-Brandis, T. Nilssen, F. Proske and T. Zhang. *A variational approach to the construction and Malliavin differentiability of strong solutions of SDE’s*. Math. Annalen, 357 (2), 761–799, 2013.
- [14] S. Peszat and J. Zabczyk. *Stochastic partial differential equations with Lévy noise*. Cambridge University Press, 2007.
- [15] E. Priola. *Pathwise uniqueness for singular SDEs driven by stable processes*. Osaka J. Math, 49, 421–447, 2012.
- [16] E. Priola. *Stochastic flow for SDEs with jumps and irregular drift term*. Stochastic analysis, 193–210, Banach Center Publ., 105, Polish Acad. Sci. Inst. Math., Warsaw, 2015.
- [17] E. Priola. *Davie’s type uniqueness for a class of SDEs with jumps*. <http://arxiv.org/pdf/1509.07448.pdf>.
- [18] E. Priola, A. Shirikyan, L. Xu, and J. Zabczyk. *Exponential ergodicity and regularity for equations with Lévy Noise*. Stoch. Proc. Appl., 122 (1), 106–133, 2012.
- [19] E. Priola, L. Xu, and J. Zabczyk. *Exponential mixing for some SPDEs with Lévy Noise*. Stochastic and Dynamics, 11, 521–534, 2011.
- [20] E. Priola, A. Shirikyan, L. Xu, and J. Zabczyk. *Exponential ergodicity and regularity for equations with Lévy Noise*. Stochastic Processes and their Applications, 122 (1), 106–133, 2012.
- [21] E. Priola and J. Zabczyk. *Structural properties of semilinear SPDEs driven by cylindrical stable processes*. Probab. Theory Related Fields, 149 (1-2), 97–137, 2011.
- [22] X. Sun and Y. Xie. *Ergodicity of stochastic dissipative equations driven by α -stable process*. Stoch. Anal. Appl., 32 (1), 61–76, 2014.

- [23] A. Ju. Veretennikov. *On the strong solutions of stochastic differential equations*. Theory Probab. Appl., 24, 354–366, 1979.
- [24] F. Y. Wang. *Gradient Estimates and Applications for SDEs in Hilbert Space with Multiplicative Noise and Dini Continuous Drift*. J. Diff. Eq., 3, 2792–2829, 2016.
- [25] F. Y. Wang. *Integrability Conditions for SDEs and Semi-Linear SPDEs*. <http://arxiv.org/pdf/1510.02183.pdf>.
- [26] L. Xu. *Ergodicity of stochastic real Ginzburg-Landau equation driven by α -stable noises*. Stoch. Proc. Appl., 123 (10), 3710–3736, 2013.
- [27] X. Zhang. *Stochastic differential equations with Sobolev drifts and driven by α -stable processes*. Ann. Inst. H. Poincaré Probab. Statist., 49, 915–931, 2013.
- [28] X. Zhang. *Strong solutions of SDEs with singular drift and Sobolev diffusion coefficients*. Stoch. Proc. Appl., 115, 1805–1818, 2005.